

## CENTERS IN LINE GRAPHS

MARTIN KNOR<sup>\*</sup>) – ĽUDOVÍT NIEPEL<sup>\*\*</sup>) – ĽUBOMÍR ŠOLTĚS<sup>\*\*\*</sup>)

(Communicated by Martin Škoviera)

**ABSTRACT.** For every graph  $H$  without isolated nodes there exists a graph  $G$  such that  $H$  is the center of  $G$  and the line graph of  $H$  is the center of the line graph of  $G$ . Graphs which are the periphery of some line graph are characterized and other distance related concepts in line graphs are studied.

### 1. Introduction and basic results

Suppose an official has to find a suitable place for an emergency facility (such as a fire station) in a given traffic network. It is naturally to locate it in such a way that the distance to the furthest node will be as short as possible, hence to build the fire station in the center of the corresponding graph. This is one reason for which centers in graphs have been studied in many papers. It is known [2, p. 41] that for each graph  $H$  there is a graph  $G$  having the center  $H$  and containing at most four noncentral nodes. The minimum number of noncentral nodes  $A(H)$  among graphs having the center  $H$  was found by Buckley, Miller and Slater [3] in the case  $H$  is a tree. Some graphs  $H$  with  $A(H) = 3$  were presented by Bielak [1] and Chen [4]. Buckley, Miller and Slater [3] have also shown that for each graph  $H$  with  $n \geq 9$  nodes and an integer  $k \geq n + 1$  there exists a  $k$ -regular graph  $G$  having the center  $H$ . So far little is known about centers of special graphs. Clearly the center of a tree consists of either a single node or a pair of adjacent nodes. All seven central subgraphs admissible in maximal outerplanar graphs were listed by Proskurowski [10]. The greatest contains six nodes. Laskar and Shier [8] studied centers in chordal graphs. The center in the cartesian product [2, p. 23] of two graphs equals the product of their centers (Niemenen [9]). Spanning subgraphs with a prescribed central node were studied by Cheston, Farley, Hedetniemi and Proskurowski [5]. They suggested an  $O(m, n)$  algorithm which for every node  $v$  in a biconnected graph  $G$  with  $n$

AMS Subject Classification (1991): Primary 05C99.

Key words: Connected graph, Center of a graph, Line graph.

nodes and  $m$  edges produces a spanning tree  $T$  such that  $v$  is a central node in  $T$ . A good survey on centers can be found in the book [2].

In this paper we focus our attention on centers in line graphs. In general, our terminology is consistent with general usage, such as in Buckley and Harary [2]. If a graph possesses no edges, then it is said to be a *trivial graph*. Let  $G$  be a connected graph,  $v$  be its node and  $f$  be its edge. Then the *eccentricity*  $e_G(v)$  or  $e(v)$  of  $v$  is the distance to a node furthest from  $v$ . The *eccentricity*  $e_G(f)$  or  $e(f)$  of the edge  $f$  equals the eccentricity  $e_{L(G)}(f)$  of the node  $f$  in the line graph of  $G$ . The *radius*  $r(G)$  is the minimal eccentricity of the nodes, whereas the *diameter*  $d(G)$  is their maximal eccentricity. Further,  $v$  is a *central node* if  $e(v) = r(G)$  and the *center*  $C(G)$  of  $G$  is the subgraph induced by all central nodes, while the *periphery*  $\text{Per}(G)$  of  $G$  is the subgraph induced by the nodes with the greatest eccentricity.

The connections in distance properties of a graph and its line graph are investigated in this paper. Relations between the eccentricity of an edge and the eccentricity of its endnodes are provided. We prove that for every graph  $H$  without isolated nodes there is a graph  $G$  such that  $H$  is the center of  $G$  and the line graph of  $H$  is the center of the line graph of  $G$ . If the line graph of a graph  $H$  has the radius at least three, then the similar result holds for the periphery instead of the center. Two conjectures on centers are presented.

We start our investigation with several observations on distances. By  $d_G(x, y)$  or  $d(x, y)$  we mean the distance between the nodes  $x$  and  $y$  in a graph  $G$ . Let  $e = ab$  and  $f = uv$  be two edges in a connected graph  $G$ . Then for their distance in the line graph of  $G$  we have  $d_{L(G)}(e, f) = 0$  if  $e = f$  and

$$d_{L(G)}(ab, uv) = 1 + \min\{d(a, u), d(a, v), d(b, u), d(b, v)\} \quad (1)$$

otherwise. Further, for an integer  $k$  and two nodes  $x$  and  $y$  in  $G$ , we mean by  $S_k(x, y)$  the subgraph in  $G$  induced by the nodes which have the distance from both  $x$  and  $y$  at least  $k$ . Now we can express the eccentricity in a line graph in the following way:

**OBSERVATION 1.** Let  $u$  and  $v$  be adjacent nodes in a connected graph  $G$  with at least three nodes. Then the eccentricity of the node  $uv$  in  $L(G)$  equals the maximal  $k \geq 0$  such that the subgraph  $S_{k-1}(u, v)$  contains an edge.

A node is *eccentric* to a node  $v$  if their distance equals  $e(v)$ . The next Lemma provides relations between the eccentricity of an edge and that of its endnodes.

**LEMMA 2.** Let  $u$  and  $v$  be adjacent nodes in a connected graph  $G$ . Then  $|e_{L(G)}(uv) - e_G(v)| \leq 1$  holds. Moreover, if  $u$  and  $v$  have distinct eccentricities,

then  $|e(u) - e(v)| =$  eccentricity of one of

**Proof.** Note that to prove that  $|e_{L(G)}(uv) - e_G(v)| \leq 1$  and Lemma 2 holds there exists an edge  $f$  such that  $e_G(v) = e(f)$ . Further, we verify that  $|e_{L(G)}(uv) - e_G(v)| \leq 1$  for  $v$  and distinct from node from  $\{u, v\}$  and there will be a  $v$  according to (1) we

A connected graph has a unique sequence of radii for the radii

**THEOREM 3.** For

(1):  $|r(L(G)) - r(G)| \leq 1$  for each two adjacent endnodes of  $f$  are and only if for each one endnode of  $f$  edge  $g$ .

(2): If  $G$  has then  $C(L(G))$  is centered, then also

(3): If  $G$  has then  $L(C(G))$  is selfcentered, then  $L(G)$  is selfcentered.

**Proof.** Part (1) assume  $G$  has the

(2): Let  $r(L(G)) = R$  and  $e(v) \geq R$ . Moreover, if  $L(G)$  in  $L(C(G))$ , so

(3): Let  $r(L(G)) = R$  and  $e(u) = e(v) = R$ , in  $C(L(G))$ . Moreover, in  $C(L(G))$ , hence

at  $v$  is a central node [2].

graphs. In general, as in Buckley and to be a *trivial graph*. its edge. Then the eccentricity  $e_{L(G)}(f)$  of the minimal eccentricity. Further, of  $G$  is the subgraph

and its line graph are ricity of an edge and at for every graph  $H$  is the center of  $G$  and . If the line graph of or result holds for the rs are presented.

distances. By  $d_G(x, y)$   $y$  in a graph  $G$ . Let  $e = f$  and

$$), d(b, v)\} \quad (1)$$

$y$  in  $G$ , we mean by ave the distance from tricity in a line graph

a connected graph  $G$  le  $uv$  in  $L(G)$  equals tains an edge.

equals  $e(v)$ . The next edge and that of its

ected graph  $G$ . Then distinct eccentricities,

then  $|e(u) - e(v)| = 1$  holds and the eccentricity of the edge  $uv$  equals the eccentricity of one its endnodes.

**Proof.** Note that  $|e(u) - e(v)| \leq 1$  as  $u$  and  $v$  are adjacent. Now it suffices to prove that  $|e_{L(G)}(uv) - e_G(v)| \leq 1$  holds. If  $G$  has two nodes, then  $G$  is  $K_2$  and Lemma 2 holds. From now on assume that  $G$  has at least two edges. Then there exists an edge distinct from  $uv$  and due to (1) we have  $e(uv) \leq 1 + e(v)$ . Further, we verify that  $e(uv) \geq e(v) - 1$  holds. Let  $a$  be a node eccentric to  $v$  and distinct from  $u$ ,  $b$  be a neighbour of  $a$ . Then the distance between any node from  $\{u, v\}$  and any node from  $\{a, b\}$  is at least  $e(v) - 2$ , since otherwise there will be a  $v - a$  path with the length shorter than  $e(v)$ . As  $uv \neq ab$ , according to (1) we have  $e(uv) \geq d(uv, ab) \geq 1 + e(v) - 2 = e(v) - 1$ .  $\square$

A connected graph is *selfcentered*, if  $C(G) = G$  holds. Now some consequences for the radius and the center in a line graph follow.

**THEOREM 3.** For a connected graph  $G$  with at least three nodes we have:

(1) :  $|r(L(G)) - r(G)| \leq 1$ , moreover,  $r(L(G)) = r(G) + 1$ , if and only if for each two adjacent central nodes  $x$  and  $y$  there is an edge  $f$  such that both endnodes of  $f$  are eccentric to both  $x$  and  $y$ . Further,  $r(L(G)) = r(G) - 1$  if and only if for each edge  $f$  joining central nodes and each other edge  $g$  at least one endnode of  $f$  has the distance at most  $r(G) - 2$  to some endnode of the edge  $g$ .

(2) : If  $G$  has a nontrivial center and a greater radius than its line graph, then  $C(L(G))$  is an induced subgraph in  $L(C(G))$ . Moreover, if  $L(G)$  is selfcentered, then also  $G$  is selfcentered.

(3) : If  $G$  has a nontrivial center and a smaller radius than its line graph, then  $L(C(G))$  is an induced subgraph in  $C(L(G))$ . Moreover, if  $G$  is selfcentered, then  $L(G)$  is selfcentered, and  $L(C(G)) = C(L(G))$  if and only if  $G$  is selfcentered.

**Proof.** Part (1) follows directly from Lemma 2 and Observation 1. Assume  $G$  has the radius  $R$ .

(2) : Let  $r(L(G)) = R - 1$ . Then for a node  $uv$  in  $C(L(G))$  we have  $e(u) \geq R$  and  $e(v) \geq R$ , Lemma 2 gives  $e(v) = e(u) = R$ , hence  $uv$  is in  $L(C(G))$ . Moreover, if  $L(G)$  is selfcentered, then  $C(L(G)) = L(G)$  is an induced subgraph in  $L(C(G))$ , so  $G$  is a subgraph in  $C(G)$  and  $G$  is selfcentered.

(3) : Let  $r(L(G)) = R + 1$ . Then, for a node  $uv$  in  $L(C(G))$ , we have  $e(u) = e(v) = R$ , which gives  $e(uv) \leq R + 1 = r(L(G))$  and that is why  $uv$  is in  $C(L(G))$ . Moreover, if  $G$  is selfcentered, then  $L(G)$  is an induced subgraph in  $C(L(G))$ , hence  $L(G)$  is selfcentered. Further, if  $G$  is selfcentered, then



we have  $L(C(G)) = L(G) = C(L(G))$ . On the other hand, suppose that  $G$  is not selfcentered. Then it contains an edge  $cy$  joining a central node  $c$  to a noncentral node  $y$ , hence  $e(c) = R$  and  $e(y) = R + 1$ . Then  $e(cy) \leq R + 1$  due to Lemma 2 and  $cy$  is in  $L(C(G))$ . Hence  $C(L(G)) = L(C(G))$  does not hold.

Theorem 4 provides results on the periphery similar to those on centers.

**THEOREM 4.** For a connected graph  $G$  with at least three nodes we have:

- (1):  $|d(L(G)) - d(G)| \leq 1$ .
- (2): If  $G$  has a nontrivial periphery and a greater diameter than its line graph, then  $L(\text{Per}(G))$  is an induced subgraph in  $\text{Per}(L(G))$ . Moreover, if  $G$  is selfcentered, then  $L(G)$  is also selfcentered and  $L(\text{Per}(G)) = \text{Per}(L(G))$  holds if and only if  $G$  is selfcentered.
- (3): If  $G$  has a nontrivial periphery and a smaller diameter than its line graph, then  $\text{Per}(L(G))$  is an induced subgraph in  $L(\text{Per}(G))$ . Moreover, if  $L(G)$  is selfcentered, then  $G$  is also selfcentered.

**Proof.** The part (1) follows directly from Lemma 2.

Denote by  $D$  the diameter of  $G$ .

(2): Let  $d(L(G)) = D - 1$ . For a node  $uv$  from  $L(\text{Per}(G))$  we have  $e(u) = e(v) = D$ , so  $e(uv) \geq D - 1 = d(L(G))$ , which gives  $uv$  is in  $\text{Per}(L(G))$ . Further, if  $G$  is selfcentered, then  $\text{Per}(G) = G$  and  $L(\text{Per}(G)) = L(G)$  is an induced subgraph in  $\text{Per}(L(G))$ , hence  $L(G) = \text{Per}(L(G))$ , which means  $L(G)$  is selfcentered.

If  $G$  is selfcentered, then  $L(G)$  is also selfcentered, and clearly,  $L(\text{Per}(G)) = \text{Per}(L(G))$  holds. On the other hand, if  $G$  is not selfcentered, then there is an edge  $py$  such that  $e(p) = D$  and  $e(y) = D - 1$ , clearly,  $py$  is not in  $L(\text{Per}(G))$ . Then, due to Lemma 2, we have  $e(py) \geq D - 1 = d(L(G))$ , hence  $py$  is in  $\text{Per}(L(G))$ , so  $L(\text{Per}(G)) = \text{Per}(L(G))$  does not hold.

(3): Let  $d(L(G)) = D + 1$ . For a node  $uv$  from  $\text{Per}(L(G))$  we have  $e(uv) = D + 1$ , so  $e(u) \geq D$  and  $e(v) \geq D$  and  $u$  and  $v$  are in  $\text{Per}(G)$ . Hence  $uv$  is in  $L(\text{Per}(G))$ . Moreover, if  $L(G)$  is selfcentered, then  $L(G) = \text{Per}(G)$  and so  $G$  is a subgraph in  $\text{Per}(G)$ , so  $G$  is selfcentered.  $\square$

## 2. Line graphs with a prescribed center

At first we show that each line graph can be a center of some line graph.

**THEOREM 5.** Let  $H$  be a connected graph  $G$  such that  $L(H)$  is the center of  $G$ .

**Proof.** Let  $V(H) = \{x_i : i = 1, \dots, n\}$ . Further the edges joining  $x_i$  to each  $x_j$  for all  $i = 1, \dots, n$  (see Fig.



Figure 1. The line

Clearly, each edge joining  $x_i$  to  $x_j$  for any other edge  $f$  let  $s$  be its distance to  $y_1z_1$  is at least  $s$ .

**CONJECTURE 6.** For a  $k$ -regular graph  $G$  there is a  $k$ -regular line graph  $L(G)$ .

Simic [11] has characterized line graphs  $L$  and the mapping  $L$ , i.e.  $L(K(G)) = K(L(G))$  for a particularly connected graph  $G$ . If  $L$  and  $C$  commute, hence  $L(C(G)) = C(L(G))$ , then, due to Theorem 3, an extra condition  $\Delta r(G) = 1$ , then, due to Theorem 4, only if  $G$  is selfcentered. Conversely, if  $G$  is selfcentered, then  $L(G) = \text{Per}(G)$  and any  $i \in \{0, -1\}$  and any  $j \in \{0, 1\}$  with  $L(C(G)) = C(L(G))$ .

**THEOREM 7.** Let  $H$  be a graph with  $n$  nodes and  $i$  be either 0 or 1. Then  $H = C(G_i)$ ,  $L(C(G_i)) = C(L(G_i))$  if and only if  $n = 4n + 6$  nodes and  $m + n^2 - 1$  edges.

hand, suppose that  $G$  has a central node  $c$  to a ... Then  $e(cy) \leq R + 1$  ...  $) = L(C(G))$  does not  $\square$

to those on centers.

three nodes we have:

diameter than its line  $G$ ). Moreover, if  $G$  is  $Per(L(G))$  holds

diameter than its line  $G$ ). Moreover, if  $L(G)$

2.

$er(G)$  we have  $e(u) = uv$  is in  $Per(L(G))$ .

$Per(G) = L(G)$  is an  $)$ , which means  $L(G)$

d clearly,  $L(Per(G)) =$  tered, then there is an  $y$  is not in  $L(Per(G))$ .  $L(G)$ , hence  $py$  is in

$(G)$  we have  $e(uv) = Per(G)$ . Hence  $uv$  is  $L(G) = Per(G)$  and so  $\square$

enter

of some line graph.

**THEOREM 5.** Let  $H$  be a graph with  $n$  nodes and  $m \geq 1$  edges. Then there is a connected graph  $G$  with at most  $4n$  nodes and at most  $m + n(n+1)$  edges such that  $L(H)$  is the center of  $L(G)$ .

**Proof.** Let  $V(H) = \{v_1, \dots, v_n\}$  be the node-set of  $H$ . Now we will construct its supergraph  $G$  as follows. Its node-set will be the set  $\{v_i, x_i, y_i, z_i \mid i = 1, \dots, n\}$ . Further the edge set consists of the edges of  $H$  together with the edges joining  $x_i$  to each node from  $V(H) - \{v_i\}$ , the edges  $x_i y_i$  and  $y_i z_i$  for all  $i = 1, \dots, n$  (see Fig. 1 for  $H = P_3$ ).




Figure 1. The line graph of the drawn graph has  $L(P_3)$  as its center.

Clearly, each edge joining two central nodes has the eccentricity three, while for any other edge  $f$  let say  $v_1$  be a central node which is nearest to  $f$ . Then its distance to  $y_1 z_1$  is at least four. Hence  $C(L(G)) = L(H)$  holds.  $\square$

**CONJECTURE 6.** For a line graph  $F$  with  $n$  nodes and each integer  $k \geq n+1$ , there is a  $k$ -regular line graph with the center  $F$ .

Simić [11] has characterized graphs  $G$ , for which the line graph transformation  $L$  and the mapping  $\mathcal{K}$  which maps a graph on its clique graph commute, i.e.  $L(\mathcal{K}(G)) = \mathcal{K}(L(G))$  holds. Now we shall study a similar class of graphs, particularly connected graphs with a nontrivial center for which the mappings  $L$  and  $C$  commute, hence  $L(C(G)) = C(L(G))$ . This is in a sense, according to Theorem 3, an extremal property. Denote  $\Delta r(G) = r(L(G)) - r(G)$ . If  $\Delta r(G) = 1$ , then, due to Theorem 3, the mappings  $L$  and  $C$  commute if and only if  $G$  is selfcentered. Complete graphs are examples of such graphs. But for any  $i \in \{0, -1\}$  and any graph  $H$  without isolated nodes, there is a graph  $G$  with  $L(C(G)) = C(L(G))$  and  $\Delta r(G) = i$ , as the next theorem states.

**THEOREM 7.** Let  $H$  be a graph with  $n$  nodes,  $m \geq 1$  edges without isolated nodes and  $i$  be either 0 or  $-1$ . Then there exist connected graphs  $G_i$  such that  $H = C(G_i)$ ,  $L(C(G_i)) = C(L(G_i))$  and  $i = \Delta r(G)$  holds. Moreover,  $G_0$  has  $4n + 6$  nodes and  $m + n^2 + 4n + 4$  edges.

Proof. Let  $V(H) = \{v_1, \dots, v_n\}$  be the node-set of  $H$  and let  $E(H)$  be its edge-set. We shall construct graphs  $G_0$  and  $G_{-1}$  with the requested properties and start with  $G_0$ .




Figure 2a. A graph  $G_0$  with  $C(G_0) = P_3$  and  $C(L(G_0)) = L(C(G_0))$ .

The node-set of  $G_0$  equals  $V(H) \cup \{a_i, b_i, c_i \mid i = 1, \dots, n\} \cup \{\alpha_1, \alpha_2, \alpha_3, \beta_1, \beta_2, \beta_3\}$ . Its edge set is  $E(H) \cup \{a_i b_i, a_i c_i, b_i c_i, \alpha_1 v_i, \beta_1 v_i \mid i = 1, \dots, n\} \cup \{c_i v_j \mid i \neq j\} \cup \{\alpha_1 \alpha_2, \alpha_2 \alpha_3, \beta_1 \beta_2, \beta_2 \beta_3\}$  (see Fig. 2a). One can check that  $r(G_0) = 3$  and  $H = C(G_0)$ , since every node outside  $H$  has the distance at least four to some of the nodes  $\alpha_3$  and  $\beta_3$ . Further, any edge from  $C(G_0)$  has the eccentricity three (its distance to  $a_1 b_1$  is three). As a center of  $L(G_0)$  lies in a single block of  $L(G_0)$ , if any edge not from  $H$  lies in  $C(L(G_0))$ , then one of its endnodes, say  $v_1$ , is in  $H$ . Then its distance to  $a_1 b_1$  is four, hence  $C(L(G_0)) = L(C(G_0))$  holds.




Figure 2b. A graph  $G_{-1}$  with  $C(G_{-1}) = P_3$  and  $C(L(G_{-1})) = L(C(G_{-1}))$ .

Now we shall construct  $G_{-1}$  with  $V(H) \cup \{a_i, b_i, c_i, d_i \mid i = 1, \dots, n\}$  (see Fig. 2b).

Obviously,  $H$  is a center of  $G_{-1}$  that every edge joining  $H$  to  $C(G_{-1})$  if an edge is adjacent to the edge  $c_1 d_1$  is at distance  $r(L(G_{-1})) = 5 = r(G_{-1})$ .

If  $H$  is bipartite, then  $C(L(G_0)) = L(C(G_0))$  property in question.

**THEOREM 8.** Let  $H$  be a graph with isolated nodes. Then  $C(L(G_0)) = L(C(G_0))$  having the center  $H$ .


Proof. Let  $A$  and  $B$  be two nodes lie in distinct blocks of  $G_0$ ,  $a, a_1, a_2, b, b_1$  and  $b_2$  are adjacent to  $b_1$  and  $b_2$  (see Fig. 3). Clearly,  $C(L(G_0)) = L(C(G_0))$ .

Nevertheless, the property  $C(L(G_0)) = L(C(G_0))$  of nodes such that  $H$  seems to be far from general.

Now we shall study the property that  $r(\text{Per}(G)) \geq d(G)$ .



t of  $H$  and let  $E(H)$   
 $-1$  with the requested



$G_0)) = L(C(G_0))$ .

$1, \dots, n\} \cup \{\alpha_1, \alpha_2, \alpha_3,$   
 $i, \beta_1 v_i \mid i = 1, \dots, n\} \cup$   
 a). One can check that  
 $H$  has the distance at  
 y edge from  $C(G_0)$  has  
 s a center of  $L(G_0)$  lies  
 lies in  $C(L(G_0))$ , then  
 to  $a_1 b_1$  is four, hence



$G_{-1}) = L(C(G_{-1}))$ .

# CENTERS IN LINE GRAPHS

Now we shall construct the graph  $G_{-1}$ . The node set of  $G_{-1}$  will contain  $V(H) \cup \{a_i, b_i, c_i, d_i \mid i = 1, \dots, n\} \cup \{x_{ij} \mid i \neq j, i = 1, \dots, n, j = 1, \dots, n\}$ . Its edge-set consists of  $E(H) \cup \{x_{ij} a_i, x_{ij} v_j \mid i \neq j\} \cup \{a_i b_i, b_i c_i, c_i d_i \mid i = 1, \dots, n\}$  (see Fig. 2b).

Obviously,  $H$  is the center of  $G_{-1}$  and  $r(G_{-1}) = 6$ , as  $d(v_i, d_i) = 6$ . Note that every edge joining central nodes has the eccentricity five due to (1). Further, if an edge is adjacent to exactly one central node, say  $v_1$ , then its distance to the edge  $c_1 d_1$  is at least six. Finally, if an edge  $f$  is adjacent to no central node, then its distance to some edge of the form  $c_i d_i$  is also at least six, so  $r(L(G_{-1})) = 5 = r(G_{-1}) + 1$  and  $L(H) = C(L(G_{-1}))$  holds.  $\square$



Figure 3. A graph having the center  $2K_2$ .

If  $H$  is bipartite, it suffices to add only six nodes in order to secure the property in question.

**THEOREM 8.** Let  $H$  be a bipartite graph on  $n$  nodes and  $m \geq 1$  edges without isolated nodes. Then there is a graph  $G$  with  $n+6$  nodes and  $m+n+4$  edges having the center  $H$  and satisfying  $C(L(G)) = L(C(G))$ .

**Proof.** Let  $A$  and  $B$  be disjoint sets of nodes in  $H$ , such that adjacent nodes lie in distinct sets. We obtain  $G$  after the addition of the new nodes  $a, a_1, a_2, b, b_1$  and  $b_2$  such that  $a$  is adjacent to  $a_1$  and to all nodes in  $A$ ,  $b$  is adjacent to  $b_1$  and to all nodes in  $B$  and  $a_2 a_1$  and  $b_1 b_2$  are also adjacent (see Fig. 3). Clearly,  $G$  has the desired property.  $\square$

Nevertheless, the problem of finding a graph  $G$  with the minimal number  $f(H)$  of nodes such that  $C(L(G)) = L(C(G))$  holds and  $G$  has a given center  $H$  seems to be far from its final solution.

## 3. The periphery in line graphs

Now we shall study the existence of line graphs with a given periphery. Note that  $r(\text{Per}(G)) \geq d(G)$  holds for each graph  $G$ .

**THEOREM 9.** *Let  $H$  be such a nontrivial graph that  $L(H)$  has the radius at most two. Then  $L(H)$  is the periphery of some line graph if and only if either  $L(H)$  is selfcentered or  $H$  contains two nodes which are not endnodes and each edge is adjacent to just one of them.*

**Proof.** If  $L(H)$  is selfcentered, then  $L(H) = \text{Per}(L(H))$  holds. Now assume  $H$  contains two nodes  $x$  and  $y$  which are not endnodes and such that each edge is adjacent to just one of the nodes  $x$  and  $y$ . Then  $L(H) = \text{Per}(L(H+xy))$  as if we add the edge  $xy$  to  $H$ , then its eccentricity will be one, while each other edge has the eccentricity two, since we have  $d_{H+xy}(xa, yb) = 2$  for pairwise distinct nodes  $a, b, x$ , and  $y$ .

Assume now that there exists a graph  $G$  such that  $L(H)$  is the periphery of  $L(G)$ . Then we have  $2 \geq r(L(H)) \geq r(\text{Per}(L(G))) \geq d(L(G))$ . Hence  $L(G)$  is either selfcentered or has the diameter two and the radius one. If  $L(G)$  is selfcentered, then  $L(H) = \text{Per}(L(G)) = L(G)$ , hence  $L(H)$  has to be selfcentered. Assume now the latter case holds. Then  $G$  contains an edge  $xy$  with the eccentricity one and so each edge is adjacent to  $x$  or  $y$ . Note that  $L(H)$  has the radius two as we have  $r(L(H)) \geq d(L(G)) = 2$ . Hence the edge  $xy$  is not in  $H$ . Further, if  $x$  is an endnode in  $H$  and  $a$  is the only its neighbour, then as  $H$  is connected there exists a node  $b$ ,  $b \neq x$  adjacent to  $ay$ . But  $b = y$  as the edge  $ab$  is adjacent to either  $x$  or  $y$  and  $a$  is distinct from  $x$  and  $y$ . Hence  $ay$  has the eccentricity one which contradicts to  $r(L(H)) = 2$ . So  $x$  and similarly  $y$  is not an endnode, which completes the proof.  $\square$

If we prescribe a line graph with the radius at least three as a periphery, then an even stronger result holds.

**THEOREM 10.** *Let the line graph of a nontrivial graph  $H$  have the radius at least three. Then  $H = \text{Per}(H + K_1)$  and  $L(H) = \text{Per}(L(H + K_1))$  holds.*

**Proof.** Let  $v$  be a node in  $H + K_1$  which is adjacent to all other nodes. As  $r(L(H)) \geq 3$  holds, we have  $r(H) \geq 2$  from Theorem 3. Hence  $v$  is the only node with the eccentricity one and each other node has the eccentricity two, so  $H = \text{Per}(H + K_1)$  holds. Further, it is easy to verify that each edge in  $H$  has the eccentricity three in  $H + K_1$  and all edges adjacent to  $v$  have the eccentricity two, that is why  $L(H) = \text{Per}(L(H + K_1))$  holds.  $\square$

Note that line g  
It would be of so  
characterized by fi  
suggested. We con  
Theorem 5. By th

# CONJECTURE 11

- [1] BIELAK, H.: *Matroids & Matroids*
- [2] BUCKLEY, F.—H. 1990.
- [3] BUCKLEY, F.—M. *a center*, J. Graph
- [4] CHEN, Z. B.: *On* (1984), 267–271.
- [5] CHESTON, G. A.—*tering a spanning t*
- [6] HEDETNIEMI, S. port CS-TR-79-11,
- [7] HEDETNIEMI, S. *C<sub>n</sub>-trees*, Utilitas M
- [8] LASKAR, R.—SHI 6 (1983), 139–147.
- [9] NIEMINEN, J.: *Th Theory*, Wishwa, G
- [10] PROSKUROWSKI, (1980), 75–79.

<sup>1</sup>Private communica



Note that line graphs are characterized by nine forbidden induced subgraphs. It would be of some interest to study centers in classes of graphs which are characterized by finite number of forbidden induced subgraphs, as Tomaszewski<sup>1</sup> suggested. We conclude with a conjecture which predicts a result analogous to Theorem 5. By the *claw* we mean the star with four nodes.

**CONJECTURE 11.** *Every claw-free graph is the center of some claw-free graph.*

## REFERENCES

- [1] BIELAK, H.: *Minimal realizations of graphs as central subgraphs*. In: *Graphs, Hypergraphs & Matroids* (Żagań, 1985), Higher College Engng., Zielona Gora, 1985, pp. 13–24.
- [2] BUCKLEY, F.—HARARY, F.: *Distance in Graphs*, Addison-Wesley, Redwood City, 1990.
- [3] BUCKLEY, F.—MILLER, Z.—SLATER, P. J.: *On graphs containing a given graph as a center*, *J. Graph Theory* 5 (1981), 427–434.
- [4] CHEN, Z. B.: *On the existence of graphs with  $A(H) = 3$* . (Chinese), *J. Math. (PRC)* 4 (1984), 267–271.
- [5] CHESTON, G. A.—FARLEY, A.—HEDETNIEMI, S. T.—PROSKUROWSKI, A.: *Centering a spanning tree of a biconnected graph*, *Inform. Process. Lett.* 32 (1989), 247–250.
- [6] HEDETNIEMI, S. M.—HEDETNIEMI, S. T.: *Centers of recursive graphs*, Technical Report CS-TR-79-11, Dept. Comp. Sc., University of Oregon (1979).
- [7] HEDETNIEMI, S. M.—HEDETNIEMI, S. T.—SLATER, P. J.: *Centers and medians of  $C_n$ -trees*, *Utilitas Math.* 21 (1982), 225–234.
- [8] LASKAR, R.—SHIER, D.: *On powers and centers of chordal graphs*, *Discrete Appl. Math.* 6 (1983), 139–147.
- [9] NIEMINEN, J.: *The center problem in the product of graphs*. In: *Recent Studies in Graph Theory*, Wishwa, Gulbarga, 1989, pp. 201–205.
- [10] PROSKUROWSKI, A.: *Centers of maximal outerplanar graphs*, *J. Graph Theory* 4 (1980), 75–79.

<sup>1</sup>Private communication, 1991.

- [11] SIMIC, S. K.: *A note on the graph equation  $C(L(G)) = L(C(G))$* , Publ. Inst. Math. (Beograd) (N.S.) **44(58)** (1988), 35–40.

Received May 8, 1991

Revised March 19, 1992

\*) Department of Mathematics

Faculty of Civil Engineering

Slovak Technical University

Radlinského 11

812 37 Bratislava

Slovakia

\*\*) Department of Applied Mathematics

Institute of Mathematic and Physics

Comenius University

842 15 Bratislava

Slovakia

\*\*\*) Department of Mathematics

Faculty of Chemical Technology

Radlinského 9

812 37 Bratislava

Slovak Technical University

Slovakia